

# Italian Journal of Agronomy Rivista di Agronomia

*An International Journal of Agroecosystem Management*

July-September 2008

Vol. 3, No. 3 supplement

10<sup>th</sup> Congress of the European  
Society for Agronomy  
15-19 September 2008, Bologna, Italy



**Multi-functional Agriculture**  
Agriculture as a Resource for Energy  
and Environmental Preservation

edited by  
Paola Rossi Pisa



The Official Journal of the  
Italian Society of Agronomy

# Italian Journal of Agronomy

## Rivista di Agronomia

*An International Journal of Agroecosystem Management*

July-September 2008 Supplement Issue

Vol. 3, No. 3 Suppl.

### Contents

<b>1. AGRICULTURE TO PRESERVE ENVIRONMENT</b>	35
<b>1.1 LAND AND WATER CONSERVATION</b>	36
<b>1.1a Erosion Control; Point and Diffuse Pollution Control (Phytodepuration, Phytoremediation and other Techniques)</b>	36
<b>Keynote Speaker: Markus Flury</b>	
<i>Soil and its Importance for Human Civilizations</i>	39
Markus Flury	
<b>Oral</b>	
<i>Application of a Crop-Farm-Indicators Modelling Chain to Assess the Impact of the EU Nitrate Directive in the Midi-Pyrenees Region</i>	41
Belhouchette H., Louhichi K., Flichman G., Therond O., Wery J.	
<i>Quantitative Parameters for the Evaluation of Cover Crop Performance in Relation to Soil Cover and Rooting Dynamics</i>	43
Bodner G., Himmelbauer M., Loiskandl W., Kaul H.P.	
<i>Water Infiltration and Soil Losses in a Permanent Bed Irrigated System in Southern Spain</i>	45
Boulal H., Gómez-Macpherson H., Gómez J.A.	
<i>Sod Seeding and Soil Erosion in a Semi-Arid Mediterranean Environment of South of Italy</i>	47
Cosentino S., Mantineo M., Copani V.	
<i>Irrigated Agriculture and Urban Waste Water: the Cases of Bean and Tomato</i>	49
D'Andrea L., Campi P., Palombo A.D., Convertini G., Ferri D., Mastroiilli M.	

<i>Land Suitability and Potential Yield Variations of Wheat and Olive Crops Determined by Climate Change in Italy</i>	797
Mereu V., Iocola I., Spano D., Murgia V., Duce P., Cesaraccio C., Tubiello F.N., Fischer G.	
<i>Modelling Yield and Yield Loss Distribution on European Scale</i>	799
Moriondo M., Bindi M., Iugeri N.	
<i>Impact of Climate Change on Grapevine (Vitis Vinifera L.) at regional Scale Phenology, Yield and Biotic Stress Responses</i>	801
Moriondo M., Bindi M., Trombi G.	
<i>Climate Change Impacts on the Guadiana River Basin: a Preliminary Study to the Adaptation Strategies</i>	803
Orioli L., Moriondo M., Brandanti G., Bindi M.	
<i>Simulating Changes of Organic Carbon Content in Soil Following Tillage Intensity and Fertilizer N Rate Reduction</i>	805
Ponzoni G., Marchetti R., Roggero P.P., Iezzi G., Seddaiu G., Corti G., Francaviglia R.	
<i>Best Practices in Agriculture: The Code of Attitudes to Prevent Mutual Impact Between Agriculture and Climate Change</i>	807
Rossi F., Di Virgilio N., Dall'Olio N.	
<i>Agro-Environmental Approach and Management of Mediterranean Archaeological Areas</i>	809
Rossi Pisa P., Bitelli G., Bittelli M., Catizone P., Ferroni L., Speranza M., Vignudelli M., Marchetti N.	
<i>Sources of Uncertainties in a Range of ETo Projections under Climate Change</i>	811
Ruiz-Ramos M., del Valle D., Mínguez M.I.	
<i>Sources of Uncertainties in Projections of Climate Change Impacts in Cropping Systems</i>	813
Ruiz-Ramos M., Mínguez M.I.	
<i>The Effect of Weather Conditions During the Growing Period on Potato Plant Development and Chlorophyll a Fluorescence Activity</i>	815
Rykaczewska K., Pietkiewicz S., Hazem Kalaji M.	
<i>Simplified Phenological Models for Open Field Tomato in Sardinia</i>	817
Salis M., Farci R., Doro L., Urracci G.R., Ledda L.	
<i>Assessing Farmers' Adaptation Capacities and Land Use Changes in Response to Forest Conservation in Madagascar</i>	819
Toillier A., Lardon S.	
<i>Management of Herbicide-Resistant Weed Beet: a Simulation Study</i>	821
Tricault Y., Sester M., Darmency H., Angevin F., Colbach N.	

# Management of Herbicide-Resistant Weed Beet: a Simulation Study

Yann Tricault<sup>1</sup>, Mathilde Sester<sup>2</sup>, Henri Darmency<sup>1</sup>, Frédérique Angevin<sup>3</sup>, Nathalie Colbach<sup>1</sup>

<sup>1</sup> UMR1210 Biologie et Gestion des Adventices, 21000 Dijon, France, colbach@dijon.inra.fr

<sup>2</sup> URP-SCRiD, CIRAD, Antsirabé, Madagascar, sester@cirad.mg

<sup>3</sup> UAR1240 Eco-innov, INRA, 78000 Thiverval-Grignon, France, Frederique.Angevin@jouy.inra.fr

In regions of sugar beet cultivation, weed beet infestations are responsible for economic losses. Weed beet belongs to the same species as the cropped plant, thus rendering herbicide control impossible in conventional sugar beet where costly practices such as manual weeding of bolters must be carried out instead. Genetically-modified herbicide-tolerant (GMHT) sugar beet varieties might provide an alternative in fields heavily infested with weed beet. However, accidental bolting of GMHT plants would result in pollen-mediated transgene flow towards weed beets. The objective of the present paper was to use a spatio-temporal simulation model for comparing 3 production systems with different cultural practices to evaluate the risk of herbicide-tolerant weed beet populations in a small agricultural region where GMHT and conventional sugar beet coexist.

## Methodology

The GENESYS-Beet model was specifically designed to evaluate different cropping systems for their potential impact on weed beet infestation and transgene dispersal through time and space (Sester et al., 2007; 2008). Model inputs are (1) the regional field pattern, (2) the crop succession in each field, (3) the cultivation techniques used to manage each crop; (4) the genotype of the sugar beet varieties (e.g. GM vs. non-GM, male sterility genes); (5) daily climate and soil conditions. The model simulates the life-cycle of weed beet in every field as a function of abiotic constraints and cropping system variables (i.e. crop, tillage operations, herbicide treatment etc.). The model operates on a daily time step with life-stage densities and genotype proportions as state variables. Genotype proportions change when the herbicide associated to the transgene is applied (i.e. glyphosate in case of GM glyphosate-tolerant sugar beet) and during pollination. Gene flow is calculated from the pollen production of flowering plants in each field and the individual pollen dispersal function integrated over the whole region for every recipient field.


Table 1. Coexisting cropping systems in farm-type simulations	Cropping system	Sugar beet variety	% regional area	Ploughing	Manual bolter- weeding	Initial weed infestation (seed/m <sup>2</sup> )
	A	GM	11	Yes	none	300
	B	non GM	45	Yes	none	0
	C	non GM	22	Yes	2	0
	D	non GM	22	No	none	0

The GENESYS-Beet simulation model was used in a case study of a French sugar beet production region where 3 types of intensive production systems had been identified (Messéan et al., 2006): (1) "potato grower", with a sugar beet/potato/winter wheat/legume/winter wheat/potato/winter wheat rotation; (2) "beet grower", with a sugar beet/winter wheat/set aside/pea/winter wheat rotation; and (3) "cereal grower", with a sugar beet/winter wheat/pea/winter wheat rotation. The three production systems were successively simulated on a real 149 field map extracted from the studied region. At the onset of each simulation, the crops of the rotation were randomly distributed in the region. In addition, four cropping systems were randomly allocated to these fields (table 1). During the subsequent years,

crops depended on crop rotation. A field once grown with a GM variety was never cultivated with a non-GM variety and vice versa. Initial seed banks were empty except for GMHT fields where infestations of weed beets justify the adoption of herbicide-tolerant varieties. Simulations lasted 15 years with the GMHT weed seed bank as the output variable.


### Results

Whatever the production system, all the fields were infested with GM seeds after 15 years. GM seed density was largest in GMHT sugar beet fields and ploughed conventional sugar beet fields without any manual weeding (Fig. 1). In unploughed and twice-weeded fields, GM seed density decreased considerably. No ploughing was even more efficient in the “cereal grower” system because the seeds were left to germinate on soil surface after sugar beet harvest, thus reducing seed survival in soil. This option is though only feasible if the following crop is a wheat where weed beet produces few seeds. Changing the management of the GM field also contributes to reducing GM seed bank in both GM and non-GM fields (Fig. 2).



### Conclusions

Our simulation results show that GM gene flow is inevitable, both in time and in space, but that it varies considerably with the cropping system. Similar results were obtained for oilseed rape. Simulation models proved to be powerful tools for predicting the effect of alternative methods on gene flow to and from weed populations. But proposals still require economic feasibility studies.



### References

Messéan A. et al. 2006. *New case studies on the coexistence of GM and non GM crops in European agriculture*. Technical Report Series of the Joint Research Center of the European Commission, EUR 22102 En, 112 p. <http://www.jrc.es/home/pages/eur22102enfinal.pdf>

Sester M. et al. 2007. Modelling the effects of cropping systems on the seed bank dynamics and emergence of weed beet. *Ecol. Model.*, 204:47–58.

Sester M. et al. 2008. GENESYS-BEET: A model of the effects of cropping systems on gene flow between sugar beet and weed beet. *Field Crop Res.*, 107:245-256.